

Press Release

July 25, 2022

FOR IMMEDIATE RELEASE

CONTACT:

Dr. Sharon Harris

516- 676-2008

safeglencove@yahoo.com

SAFE Glen Cove Coalition: NIDA Percentage of Methadone Over Dose Deaths Decline

According to a new study by the National Institute on Drug Abuse (NIDA), the percentage of methadone-involved overdose deaths relative to all drug overdose deaths declined from January 2019 to August 2021. Access to methadone, a medication to treat opioid use disorder, was expanded at the start of the COVID-19 pandemic to allow more patients to take home doses, rather than visit a clinic daily. These data indicate that broader access to treatment was not associated with harms. While drug overdose deaths both with and without methadone increased in the month of March 2020, overdose deaths that did not involve methadone continued to increase in the months after the policy changes, while overdose deaths involving methadone held steady. This study was a collaborative effort led by researchers at NIDA, part of the National Institutes of Health, and the National Center for Injury Prevention and Control, part of the Centers for Disease Control and Prevention.

In 2021, provisional data from CDC estimate more than 107,000 people died of a drug overdose, with 75% of those deaths involving an opioid. The overall rise in overdose deaths is largely attributable to the proliferation in the drug supply of illicit fentanyl, a highly potent synthetic opioid. A key component of the federal government response to the overdose crisis is expanding access to medications for opioid use disorder. However, only 18% of people with opioid use disorder receive medication as treatment. Though the benefits of providing safe and effective medication for opioid use disorder are well-known, decades of stigma against treating substance use disorders with medication has contributed to minimal reach.

In the United States, methadone for the treatment for opioid use disorder can only be provided through federally certified opioid treatment programs, where most patients are required to visit a clinic in-person, on a daily basis, in order to get their medication. For decades, this requirement has been identified as an often-insurmountable barrier to access and retention for this treatment, particularly for people trying to balance employment, childcare, and other needs. The requirement presented unique challenges during the COVID-19 pandemic, as accessing in-person treatment became limited due to concerns about exposure to COVID-19. In order to ensure continuity of care for individuals receiving methadone treatment, on March 16, 2020, the Substance Abuse and Mental Health Services

Administration (SAMHSA) allowed states to request exceptions to provide up to 28 days and 14 days of take-home methadone for stable and less stable patients, respectively.

To assess the impact and potential harms of these policy changes, investigators used data from January 2019 through August 2021 from the CDC's National Vital Statistics System, a national mortality database. Researchers calculated monthly drug overdose deaths without methadone, monthly drug overdose deaths involving methadone, and the percentage of overall overdose deaths involving methadone. They then assessed whether there was a shift in outcomes before and after the methadone take-home policy change in March 2020. They did this through interrupted time series analyses, a method of evaluation for large scale public health interventions with well-defined starting points.

Researchers found that non-methadone-involved overdose deaths increased by an average of 78 more deaths each month before March 2020, increased by 1,078 deaths during March 2020, and then continued to increase by an average of 69 more deaths each month after March 2020. Methadone-involved overdose deaths experienced a similar increase in March 2020 (increased by 94 deaths). However, the trend in number of deaths per month before and after this initial uptick remained stable, and the percentage of overdose deaths involving methadone declined at similar rates before and after the take-home policy change, declining from 4.5% of overdose deaths in January of 2019 to 3.2% in August 2021.

Taken together, these findings indicate that the modest increase in methadone-involved overdose deaths in March 2020 was likely a reflection of the overall spike in overdose deaths driven by illicitly produced fentanyl and not an outcome of the take-home policy change. Mirroring findings from smaller studies, these national data provide evidence that the expanded opioid treatment program take-home methadone policy change established in March 2020 was not associated with increases in methadone-involved overdose deaths, despite marked increases in overall overdose deaths during the study period. Coupled with other studies that have demonstrated positive benefits related to these policies, the authors note that these findings can inform decisions about permanently expanding take-home methadone from opioid treatment programs.

NIDA researchers maintain this evidence adds significant weight to the argument that effective treatment for substance use disorders should be offered in an accessible and practical way that works for people who need it. The goal of health policy should be to promote health and reduce harm, and conducting research studies like this ensures that those policies are based on the best available scientific evidence.

NIDA is a component of the National Institutes of Health, U.S. Department of Health and Human Services and supports most of the world's research on the health aspects of drug use and addiction. For more information about NIDA and its programs, visit www.nida.nih.gov.

SAFE is the only alcohol and substance use prevention agency in Glen Cove. Its Coalition is conducting an opioid prevention awareness campaign entitled, "Keeping Glen Cove SAFE," in order to educate and update the community regarding opioid use and its consequences. To learn more about the SAFE Glen

Cove Coalition please follow us on www.facebook.com/safeglencovecoalition or visit SAFE's website to learn more about the Opioid Epidemic at www.safeglencove.org.